

Anti-Nausea Medication Guide with Aprepitant (8 Day Regimen)






Patient Name: _____

Chemo Regimen: _____

Cycle Number: _____

Instructions for patient:





Please refill your anti-nausea medications **BEFORE** your next cycle. Bring your anti-nausea medications on your treatment day.

Note: Images are not to scale	Day 1 : Treatment Day		Day 2		Day 3		Day 4		Day 5		Day 6		Day 7		Day 8	
	*Before chemo	After chemo (PM)	*Before chemo	After chemo (PM)	*Before chemo	After chemo (PM)	*Before chemo	After chemo (PM)	*Before chemo	After chemo (PM)	AM	PM	AM	PM	AM	PM
Aprepitant (Emend®) 125 mg White & pink Capsule  Take with or without food	Take 1 125 mg capsule															
Aprepitant (Emend®) 80mg White capsule  Take with or without food			Take 1 80 mg capsule		Take 1 80 mg capsule		Take 1 80 mg capsule		Take 1 80 mg capsule		Take 1 80 mg capsule		Take 1 80 mg capsule			
Ondansetron (Zofran®) 8 mg Yellow, oval  Take with or without food	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet	Take 1 8mg tablet						
Dexamethasone (Decadron®) 4 mg White, 5-sided  Take with food	Take 3 tablets (12 mg total)		Take 2 tablets (8 mg total)								Take 1 4mg tablet	Take 1 4mg tablet	Take 1 4mg tablet	Take 1 4mg tablet	Take 1 4mg tablet	Take 1 4mg tablet
Prochlorperazine 10 mg Orange, round  Take with or without food	<ul style="list-style-type: none"> You can take 1 tablet every 4 to 6 hours when needed for nausea or vomiting (up to a maximum of 4 tablets (40mg) per day) You can start taking this tablet on “Day 1: Treatment Day” and continue as needed when nauseated 															

*Before chemo: 30-60 minutes before your chemo appointment.


* After Chemo: 12 hours after your “before chemo” dose

Your Medications, Common Side Effects & How to Manage Them:

	<p>Aprepitant (Emend®) prevents nausea and vomiting. Common side effects can include diarrhea, constipation, upset stomach, dizziness, hiccups, and fatigue. If you have trouble breathing, swelling, and/or a rash appears, stop taking the drug and let us know about these symptoms.</p>
	<p>Ondansetron (Zofran®) works well to prevent nausea and vomiting after you have received your chemotherapy. Common side effects can include constipation, and/or headache.</p>
	<p>Dexamethasone (Decadron®) works well to help control your nausea but it may increase your blood sugar if you have diabetes, interfere with sleep, increase appetite, cause nervousness, or cause moodiness.</p> <p>If you have diabetes, check your blood sugars regularly and work with your family doctor to help keep blood sugars near normal. Speak with your oncology nurse or doctor to manage other side effects.</p>
	<p>Prochlorperazine is used to help prevent nausea and vomiting following chemotherapy. Common side effects can include drowsiness, dry mouth, and/or constipation.</p> <p>If you experience any difficulty with emptying your bladder or any new muscle twitching or jerking, please stop the medication and let your oncology nurse or doctor know about this problem.</p>

Other tips:

- Drink at least 2 litres of non-caffeinated fluids per day, unless told otherwise by your cancer care team. Staying hydrated helps to prevent nausea.
- Use *My Health Diary* to track your symptoms and bring the Diary each time you come
- Complete your symptom screening to rate your symptoms at each doctor visit
- Before you start your anti-nausea medications, tell your cancer care team all the medications you're taking including any over-the-counter



**If you have questions about your medications call the Cancer Symptom Support Telephone Service:
 During the day: (705) 728 – 9090 ext. 79565. Evenings, weekends and holidays: 1-877-681-3057.
 Please visit www.rvh.on.ca for the latest hours of operation.
 Bring this sheet and your medications to your next chemotherapy treatment appointment.**