



GROUP A *STREPTOCOCCUS* DISEASE

CoP MEETING - RVH IPAC HUB

Mustansir Diwan MSc, CIC, LTC-CIP – February 16, 2023


WHAT IS GAS AND INVASIVE GAS?

- Group A *Streptococcus* (GAS) are bacteria commonly found in the throat and on the skin.
- People can be carriers of these bacteria with no symptoms of illness.
- Spreads from person-to-person through direct or indirect **contact** with secretions from nose, throat, wound secretions or respiratory **droplets**.
- Incubation period of group A strep infections is usually 1-3 days.
- Period of communicability could be from weeks or months when **not on** antibiotics.

WHAT IS GAS AND INVASIVE GAS? (CONT.)

- Common non-invasive GAS infections include Strep throat, scarlet fever, impetigo, and ear infections.
- In rare cases, infection can become “**invasive**” when bacteria enter sterile parts of the body such as blood, deep tissue or the lining of the brain.

THE PATHOGEN



Source: www.osmosis.org

STREPTOCOCCUS PYOGENES




Image source: phil.cdc.gov

HOW COMMON IS iGAS DISEASE?

- A total of 329 confirmed cases of iGAS reported in Ontario for the 2022-2023 iGAS season as of January 23, 2023.
- Highest incidence reported among those ≥ 65 years of age.
- In Ontario, the incidence rate is 2.1 iGAS cases per 100,000 population.
- In Canada, the incidence of iGAS disease increased from 4.0 to 8.1 cases per 100,000 from 2009 to 2019.
- iGAS disease remains an important public health concern.

WHO IS AT RISK?

- Individuals with weaker immune system:
 - Infants and elderly
 - HIV infection
 - Diabetes
 - Malignancy
 - Underlying medical condition
- Bacteria can get into the blood stream usually through a breach in the skin, mucosal laceration, or following surgery.

SYMPTOMS

- Pharyngitis: fever, painful swallowing, tonsils look reddish and swollen.
- Impetigo: itchy, honey-coloured scabs on the skin.
- Erysipelas: warm, painful lesions on the skin with raised edges.
- Scarlet fever: fever, bright-red skin rash, tongue with sandpaper feel.
- Necrotizing Fasciitis: fever with affected skin becoming painful and purple in colour – invasive GAS*
- Streptococcal Toxic Shock Syndrome (STSS): fever, chills, myalgia, nausea, vomiting – invasive GAS*

DIAGNOSIS

- In Canada, iGAS is **reportable** in all provinces and territories. Confirmed cases are notifiable at the national level.
- Rapid Strep Test (RST) or culture from throat swabs usually used for non-invasive GAS infection.
- For skin infections, cultures can be done from blisters or debrided tissues.

CASE DEFINITION FOR INVASIVE GAS

- **Confirmed case:** laboratory confirmation of infection with or without clinical evidence of invasive disease. Requires isolation of group A streptococcus (*Streptococcus pyogenes*) from a normally sterile site like blood, cerebrospinal fluid, joint, pleural, pericardial fluid.

OR

- isolation of group A streptococcus from a non-sterile site (e.g., skin) with clinical evidence of severity.

CLOSE CONTACT DEFINITION IN LTCH AND RH

- Individual must have exposure to the case during the period from 7 days prior to onset of symptoms in the case to 24 hours after the case's initiation of antimicrobial therapy.
- Spending ~4 hours per day on average in the previous 7 days or 20 hours per week with the case.
- Those with following types of contact with the iGAS case:
 - Sharing the same bed, sexual relations, direct mucous membrane contact with oral or nasal secretions, or direct contact with the open skin lesion, shared needles.

CASE MANAGEMENT

- Report confirmed case and close contacts immediately to the local public health unit.
- Conduct a retrospective chart review of all residents/clients over the previous 4 to 6 weeks:
 - For culture-confirmed cases of GAS disease
 - For any suggested cases of non-invasive or invasive GAS infection
 - Exclude pneumonia and conjunctivitis not confirmed by culture
- Assess the potential for a source of infection from outside the home.

IPAC MEASURES FOR iGAS DISEASE

- Symptomatic residents/clients should be placed on **Contact and Droplet Precautions** immediately until 24 hrs after initiation of ABT.
- Have staff wear appropriate PPE when providing care to resident on isolation.
- Practice good hand hygiene and other routine practices at all times.
- Do not allow symptomatic residents to participate in group activities for at least 48 hrs after symptoms have resolved.
- Use dedicated equipment for residents on isolation when possible.
Thoroughly clean and disinfect shared items after each use.

TREATMENT



REFERENCES

- Golden AR, Griffith A, Demczuk WHB, Tyrrell GJ, Kus JV, McGeer A, Domingo M-C, Hoang L, Minion J, Van Caeseele P, Smadi H, Haldane D, Zahariadis G, Mead K, Steven L, Strudwick L, Li AY, Mulvey MR, Martin I. Invasive group A streptococcal disease surveillance in Canada, 2020. *CCDR* 2022; 48(9):407-14.
<https://doi.org/10.14745/ccdr.v48i09a05>.
- Ontario Agency for Health Protection and Promotion (Public Health Ontario), *Provincial Infectious Diseases Advisory Committee*. Recommendations on Public Health Management of Invasive Group A Streptococcal (iGAS) Disease in Ontario. Toronto, ON: Queen's Printer for Ontario; 2014.
- Ontario Agency for Health Protection and Promotion (*Public Health Ontario*). Invasive Group A streptococcal disease in Ontario: 2016-17 seasonal summary. Toronto, ON, Queen's Printer for Ontario; 2018.
- Ontario Agency for Health Protection and Promotion (*Public Health Ontario*). iGAS in children 0 to 17 years of age in Ontario: October 1, 2016 to January 23, 2022. Toronto, ON: King's Printer for Ontario; 2023.
- Public Health Agency of Canada. Guidelines for the Prevention and Control of Invasive Group A Streptococcal Disease. *CCDR* 2006; 32S2:1-26.

REFERENCES (CONT.)

- Simcoe Muskoka District Health Unit. Invasive group A streptococcal infections (iGAS), *SMDHU*; 2017. Available at: <https://www.simcoemuskokahealth.org/Topics/InfectiousDiseases/DiseaseInformation/FactSheetsIL/IGAS.aspx>. Accessed February 08, 2023.
- Public Health Agency of Canada. Pathogen Safety Data Sheet: Infectious Substances - Streptococcus pyogenes. *Government of Canada*; 2010. Available at: <https://www.canada.ca/en/public-health/services/laboratory-biosafety-biosecurity/pathogen-safety-data-sheets-risk-assessment/streptococcus-pyogenes.html>. Accessed February 15, 2023.
- Simcoe Muskoka District Health Unit. Group A Streptococcal disease, invasive. *SMDHU*; 2017. Available at: <https://www.simcoemuskokahealth.org/docs/default-source/jfy-health-care-professionals/reportable-diseases/group-a-strep.pdf>. Accessed February 15, 2023.
- Appendix 1: Case Definitions and Disease-Specific Information - Disease: Group A Streptococcal Disease, invasive (iGAS). 2022. Ontario Public Health Standards - Infectious Disease Protocol. *Ministry of Health*. Available at: https://www.health.gov.on.ca/en/pro/programs/publichealth/oph_standards/docs/gas_chapter.pdf. Accessed February 15, 2023.

QUESTIONS?

